

EXCEPTIONAL ALGEBRAIC RELATIONS FOR RECIPROCAL SUMS OF FIBONACCI AND LUCAS NUMBERS

Carsten Elsner, Shun Shimomura, and Iekata Shiokawa

**Diophantine Analysis and Related Fields 2011, AIP Conf. Proc., no. 1385,
M. Amou and M. Katsurada (Eds.), A.I.P., 2011, 17-31**

Let $(F_n)_{n \geq 0}$ and $(L_n)_{n \geq 0}$ be the sequences of Fibonacci and Lucas numbers, respectively. Moreover, we define a set Γ of 12 real numbers by

$$\Gamma := \left\{ \sum_{n=1}^{\infty} \frac{1}{F_n^{2s}}, \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{F_n^{2s}}, \sum_{n=1}^{\infty} \frac{1}{L_n^{2s}}, \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{L_n^{2s}} \quad (s = 1, 2, 3) \right\}.$$

From former investigations of the authors it is known that every 4-subset of Γ consisting of four numbers is algebraically dependent over \mathbb{Q} . It is also proven by the authors that each of the 3-subsets

$$\left\{ \sum_{n=1}^{\infty} \frac{1}{F_n^{2s}} \quad (s = 1, 2, 3) \right\}, \left\{ \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{F_n^{2s}} \quad (s = 1, 2, 3) \right\}, \left\{ \sum_{n=1}^{\infty} \frac{1}{L_n^{2s}} \quad (s = 1, 2, 3) \right\},$$

$$\left\{ \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{L_n^{2s}} \quad (s = 1, 2, 3) \right\}$$

is algebraically independent over \mathbb{Q} . There are

$$\binom{12}{3} = 220$$

3-subsets of Γ . In this paper it is shown that among them there are exactly 22 such 3-subsets which are algebraically dependent over \mathbb{Q} . We compute the corresponding polynomials explicitly. For instance, we have

$$2 \sum_{n=1}^{\infty} \frac{1}{F_n^2} - \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{F_n^2} - 5 \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{L_n^2} = 0,$$

$$\left(\sum_{n=1}^{\infty} \frac{1}{F_n^2} \right) \cdot \left(8 \sum_{n=1}^{\infty} \frac{1}{L_n^2} + 1 \right) - 5 \sum_{n=1}^{\infty} \frac{1}{L_n^2} - 20 \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{L_n^4} = 0.$$

MR 2010 Subject Classification: 11J81, 11J91, 33E05

Key words: Fibonacci numbers, Lucas numbers, reciprocal sums